

## Potentiality of *Moringa oleifera* aqueous extract as a growth modulator and antistress in acute hypoxic Nile tilapia *Oreochromis niloticus*

Shourbela R.M.<sup>1</sup>; El-Hawarry W.N.<sup>1\*</sup>; Abd El-Latif A.M.<sup>2</sup>; Abo-Kora S.Y.<sup>3</sup>

Received: April 2017

Accepted: December 2017

### Abstract

This study aims to get a comprehensive evaluation of the growth promoting effects and the hypoxic stress relief potentiality of *Moringa oleifera* aqueous extracts. *Oreochromis niloticus* fingerlings were arbitrarily allocated into five duplicated fish groups (30 fish tank<sup>-1</sup>). The fish groups were labeled according to the *M. oleifera* aqueous extract dietary inclusion level (G<sub>1</sub>:G<sub>5</sub>). MOAE had fundamentally promoted tilapia growth. Serum total protein levels were considerably higher in the *M. oleifera* fed fish, whereas the levels of liver enzymes diminished significantly in G<sub>5</sub> fish. Additionally, the dietary *M. oleifera* resulted in a noticeable hypoglycemic effect together with a pronounced decline in the antioxidant activities. The use of *M. oleifera* supplemented diet decreased the hypoxia-related stress as conveyed by the gradual descent in the serum cortisol levels of the hypoxic-stressed tilapia. This study proposes the potentiality of *M. oleifera* aqueous extract as a growth promoter, antistress and antioxidants. It also validates its safe application in commercial tilapia culture. Future study is required to comprehend the influence of this plant extract in relieving chronic stress and its possible toxic effect as well. Feasibility study for its commercial usage is required too.

**Keywords:** Plant extracts, *Moringa oleifera*, *Oreochromis niloticus*, Growth, Antioxidant activity, Hypoxic stress.

1-Department of Animal Husbandry and Animal Wealth Development, Faculty of Veterinary Medicine, Alexandria University, Edfina, 22758, Rosetta line, Egypt.

2-Department of Fish Diseases and Management, Faculty of Veterinary Medicine, Benha University, Egypt.

3-Department of Pharmacology, Faculty of Veterinary Medicine, Benha University, Egypt.

\*Corresponding author's Email: waleed.elhawarry@alexu.edu.eg

## Introduction

Tilapia is a popular aquatic species being cultured worldwide. In 2012, the world tilapia production had boomed to reach 4.5 MT (FAO, 2014) and is anticipated to increase exponentially over the coming years. Indeed, Tilapia intensification is an essential requisite to cover the existent necessity for fish protein. The generated stress and limited disease control associated with intensive aquaculture have made sound management and biosecurity critical challenges to aquaculture (Kautsky *et al.*, 2000; Schreck, 2000; Moss *et al.*, 2012). As an important water quality criterion; Dissolved oxygen (DO) may represent a significant constraint for successful aquaculture. For instance oxygen depletion may occur under different culture conditions especially in intensive culture systems prompting fish hypoxia. Hypoxia has many adversative consequences on cultured fish. It can antagonistically affect varied kind of physiological and biochemical activities including growth (Wang *et al.*, 2009) and hematological indices along with stress response stimulation (Simi *et al.*, 2016).

Antimicrobials and other chemicals are regularly administered as additives in fish diets or water baths as growth promoters, prophylactics or therapeutants (Bulfon *et al.*, 2015). Their continuous application resulted in various adversarial effects for the environment and health safety (Menz *et al.*, 2015). Accordingly, numerous countries have rigid regulations that limit their application in aquaculture. In this concern, a new worldwide trend

had evolved regarding the consumer perceptions as influenced by safety and quality of the animal crop. In this manner, imperative needs to search different choices to antimicrobials were of primary concerns. Such choices would guarantee better growth performance and pathogens control and afterward, would polish the sustainable fish production under intensive culture systems. Plants are the warehouses of safe and inexpensive chemicals. Natural plants may represent better substitutes for antimicrobials in aquaculture as they impound various actions like growth enhancement, antimicrobial and antioxidant activities together with their antistress potentialities (Reverter *et al.*, 2014). Indeed, this is related to the several active constituents they have, such as pure volatile oils, alkaloids, phenolics, terpenoids, saponins, flavonoids and numerous other compounds (Anwar *et al.*, 2007; Chakraborty and Hancz, 2011; Huyghebaert *et al.*, 2011).

The drumstick tree, *Moringa oleifera*, is angiosperm plant demonstrating varied and valuable impacts, in accord to the plant part and inception. Different portions of this plant have been used all through history for its nourishing and healing importance (Mbikay, 2012; Leone *et al.*, 2015). The seeds, for instance, possess potential antimicrobial action against certain pathogens. They also can be utilized for water cleansing since they contain particular proteins with coagulation properties (Nikkon *et al.*, 2003; Suarez *et al.*, 2003). Nevertheless, the information available

about its influences on the immune response, antioxidant activity and growth of various fish species including Nile tilapia *Oreochromis niloticus*. To the best of our information, there is no data in the literature on the influence of dietary *M. oleifera*, in evoking a better fish response to stressful culture conditions, such as hypoxia. Therefore; this study aims to get a comprehensive evaluation of growth promoting effect of *M. oleifera* aqueous extracts (MOAE) on *O. niloticus* fingerlings. In addition, the MOAE influences on the enzymes antioxidant activities and some serum biochemical indices of acutely hypoxic Nile tilapia fingerlings.

## Materials and methods

### *Fish and rearing*

Apparently, healthy monosex tilapia fingerlings obtained from a private fish hatchery at Kafr El Sheikh Governorate, Egypt. The fish were transferred into polyethylene bags filled with oxygen to the Lab of Fish Breeding and production, Vet. Med, Alex. University, Egypt. Fingerlings were firstly acclimated in rectangular white plastic tanks (500 L capacity) supplied with 300 L of underground freshwater and equipped with individual biofilters along with constant air supply. During the adaptation period, fish were fed twice daily (09:00; 14:00) with a diet previously formulated to obtain a 30% crude protein.

### *Preparation of aqueous extract of M. oleifera leaves*

Mature leaves of *M. oleifera* were

selected and gathered from one area (Abu-hammad, Sharkia, Egypt). The air dried leaves were crushed using mortar and pestle. The Bioactive substances of *M. oleifera* leaves were extracted using infusion technique. The leaves were immersed in distilled water for 24 h using 1:2 ratios (weight /volume) (Fernandez, 1990). After that, a suction apparatus and doubled Whatman no. 1 filter paper used to spate the debris and filtrate. Then a concentrated filtrate was obtained through a vacuum rotary evaporator (Buchi R-110 Rotavapor, Switzerland) at (40°C) the dry extract kept frozen at 0°C.

### *Diet preparation*

Five diets were prepared (Table 1), include the control one which was non-*M. oleifera* supplemented diet and the other four ones were supplemented with graded levels (50, 100, 200 and 400 mg Kg<sup>-1</sup> diet) of *M. oleifera* watery extract. Then, tap water was added to each until a firm paste was acquired. Each doughy diet was then separately passed through a mincer with a 16 mm die, the resulted strands were gently broken into pellets, air dried at ambient temperature for two days and finally kept at 4°C in plastic bags till need.

### *Experimental setup and fish management*

After adaptation periods, fish with an average body weight of 4.48±0.22 g and average body length of 6.27±0.19 cm were arbitrarily allocated into five replicated fish groups (two replicates group<sup>-1</sup>). The fish groups were differentiated according to the MOAE

dietary inclusion level. They were labeled as G<sub>1</sub>; fed a control diet (0% MOAE kg<sup>-1</sup> feed), G<sub>2</sub>; fed 50 mg MOAE kg<sup>-1</sup> feed, G<sub>3</sub>; fed 100 mg MOAE kg<sup>-1</sup> feed, G<sub>4</sub> was fed 200 mg MOAE kg<sup>-1</sup> feed and G<sub>5</sub>; fed 400 mg MOAE kg<sup>-1</sup> feed. The stocking density was maintained at 30 fish tank<sup>-1</sup>. One day after stocking, each fish group was hand-fed with its corresponding diet. The trial was continued for 60 days, with a feeding frequency two times per

day (09:00; 14:00), six days a week until apparent satiation. Likewise, the adaptation period, the water quality of each tank was similarly preserved and managed throughout the experimental period. Moreover, 1/3 of the water in all tanks was replaced each other day. Accordingly, the tanks water quality was within the permissible limits (temperature; 26±0.42 °C, pH; 7.87±0.36, and DO; 6.94±0.56 mg L<sup>-1</sup>).

**Table 1: Composition and proximate analysis of the basal diet (g 100 g<sup>-1</sup> dry matter).**

Feed ingredients	Proportion
Fish meal (60%)	16
Soybean meal (47%)	27.5
Yellow corn	27
Rice bran	13
Wheat bran	15
Dicalcium phoshate	1
Methionin	0.05
Cholin chloride	0.05
Vitamin and mineral premix*	0.2
Binder	0.2
Total	100
<b>Composition</b>	<b>Proximate analysis (%)</b>
Dry matter	88.09
Crude protein	29.33
Crude Fiber	5.58
Ash	8.15

\*Vitamin mineral premic (Technomix) (quantity per 1 kg).  
 Vitamin premix: Vitamin-A: 12,000,000 IU; Vitamin-D3: 2,000,000 IU; Vitamin-E: 10,000 mg; Vitamin-K<sub>3</sub>: 2,000 mg; Vitamin-B<sub>1</sub>: 1,000 mg; Vitamin-B<sub>2</sub>: 5,000 mg; Vitamin-B<sub>6</sub>: 15,000 mg; Vitamin-B<sub>12</sub>: 1.0 mg; Biotin: 50 mg; Pantothenate: 10,000 mg; Nicotinic acid: 30,000 mg; Folic acid: 1,000 mg.  
 Mineral premix (mg per 2 kg): FeSO<sub>4</sub>: 30,000; ZnSO<sub>4</sub>: 50,000; MnSO<sub>4</sub>: 60,000; CuSO<sub>4</sub>: 10,000; calcium iodine: 1,000; cobalt: 100; choline: 250,000.

*Fish growth and survival*

The body weight and the total length of all experimental fish groups were noted 24 h after the end of the feeding trial. The performance data and the survival rate (%) were appraised;

- Weight gain (g) =  $W_2 - W_1$
- Average daily gain (g day<sup>-1</sup>) =  $(W_2 - W_1) / T$
- Specific growth rate (% day<sup>-1</sup>) =  $[\log_e (W_2) - \log_e (W_1) / T] \times 100$
- Condition factor (K) =  $(W/L^3) \times 100$

Where;  $W_2$  is final weight (g),  $W_1$  is initial weight (g), T is the trial period (days), L is total length (cm).

Hepatosomatic index (%) =  $[\text{liver weight} / W] \times 100$

Survival rate (%) =  $[\text{Number of survived fish} / \text{initial number of fish}] \times 100$

*Biochemical parameters*

To evaluate some of the biochemical parameters, blood samples (three anesthetized fish tank<sup>-1</sup>) were collected from either the heart or the caudal vessels 24 h after the last diet. Hypodermic syringes were used to get the blood samples which were then transmitted to Wassermann tubes. The serum samples were obtained through allowing blood clotting at room temperature for 45 min then centrifuged at 3000 rpm (Hettich Centrifuge, Tuttlingen, Germany) for 15 minutes (Burnett *et al.*, 2011). The sera were pipetted into Eppendorf tubes, labeled and hold on in deep freeze at -20 °C for further biochemical analysis.

Biochemical indices such as total protein and albumin were measured in

step with Lowry *et al.* (1951) and Drupt *et al.* (1974). Immunoglobulin M (IgM) was assessed via spectrophotometric examination of fish serum. Also, the hepatic transaminase activities (ALT and AST) were tested in serum based on the method displayed by Reitman and Frankel (1957). Whereas serum urea levels were analyzed according to Fawcett and Scott (1960) and creatinine levels were estimated after Husdan and Raporpot (1968). Serum glucose levels were evaluated by the methodology of Cooper and Mc Daniel (1970).

*Antioxidant enzymes assay*

From each replicates three randomly selected fish were seined and transferred directly into an anesthetic containing water. Each fish was then dissected and the liver was extracted, cleaned by 0.9% NaCl solution. After that, each liver was homogenized in a cooled phosphate buffer saline (pH 7.2 at a ratio 1: 10) using electro-homogenizer (Heidolph, Germany). The homogenate was then centrifuged (13,000 ×g at 4°C for 10 min) and the supernatant was pipetted and stored at -80°C until analysis. The antioxidant enzymes assay comprised; superoxide dismutase (SOD), Glutathione peroxidase (GPX), (Glucose 6-phosphate dehydrogenase (G6PDH) and Nitric oxide (NO) which were analysed according to Paglia and Valentine (1967), Nishikimi *et al.* (1972), Bautista *et al.* (1988) and Van Bezooijen *et al.* (1998); respectively.

*Acute hypoxic stress*

Toward the completion of this trial,

tilapia from various treated tanks were exposed to acute hypoxic stress. The tilapia fish were retained out of their tank and subjected to atmospheric air for three minutes. Investigation of cortisol profile was carried out through taking blood samples (5 fish treatment<sup>-1</sup>). Thirty minutes and again two hours later from the same fish were sourced out of water for blood sampling to detect the post-hypoxia cortisol profile (Knobil, 1980).

#### Statistical analysis

One-way ANOVA was used to statistically analyze the data recorded for the differently treated fish groups under SAS (2008). Duncan's multiple range tests applied as well to detect any anticipated significant differences between treated fish groups at a significant level of 95%.

#### Results

Feeding MOAE supplemented diet had significantly enhanced tilapia growth and survivability in comparison to the MOAE non-supplemented one (Table 2). The FBW, WG, ADG, SGR and FL had significantly increased ( $p < 0.05$ ) in an ascending trend as the dietary inclusion level of MOAE increased from 50 mg kg<sup>-1</sup> diet to 400 mg kg<sup>-1</sup> diet. The condition factor was not significantly ( $p > 0.05$ ) differed by the MOAE dietary supplementation. However, higher estimates were recorded for the MOAE received fish. The MOAE supplemented diet expressively ( $p < 0.05$ ) increased the HSI, where the highest indices were assessed at 400 mg MOAE kg<sup>-1</sup> diet

inclusion level. Fish fed MOAE supplemented ration expressed significantly ( $p < 0.05$ ) better FCR values. Additionally, the lowest FCR values were achieved by those fish fed diets in 100 and 400 mg MOAE kg<sup>-1</sup>, respectively. The Survival percentage noted in this study was not ( $p > 0.05$ ) affected by the MOAE diet supplementation.

Dietary MOAE significantly ( $p < 0.05$ ) influenced the serum TP level of *O. niloticus* fish (Table 3). The addition of 400 mg MOAE kg<sup>-1</sup> diet resulted in an expressively higher serum TP level than the levels identified in the other treated fish groups. However, still comparable results were observed between the control MOAE non-supplemented group and the other treated fish groups (G2, G3, and G4). Meanwhile, albumin levels were not considerably ( $p > 0.05$ ) influenced by MOAE dietary supplementation. Additionally, IgM values noticeably increased ( $p < 0.05$ ) in the fish supplemented with 200 mg MOAE kg<sup>-1</sup> diet. Liver and kidneys enzymes are demonstrated in Table 3; the G<sub>5</sub> exposed a noteworthy ( $p < 0.05$ ) drop in liver enzymes. The recorded urea levels diminished significantly in the fish fed 100 mg MOAE kg<sup>-1</sup> diet. However, the serum creatinine levels did not elucidate any substantial differences ( $p > 0.05$ ) as an influence to dietary MOAE supplementation. MOAE considerably reduced the glucose levels. ( $p < 0.05$ ) in the sera of the MOAE supplemented fish (Table 3). Both G4 (200 mg kg<sup>-1</sup>) and G1 (control one) displayed the highest

serum glucose concentrations.

**Table 2: Growth performance of *Oreochromis niloticus* fed dietary *Moringa oleifera* aqueous extract supplementation for 60 days.**

Parameters	Treatments				
	Dietary <i>M. oleifera</i> extract mg kg <sup>-1</sup> diet				
	0 (Control)	50	100	200	400
IBW(g)	4.45±0.24	4.53±0.23	4.53± 0.24	4.48±0.23	4.40±0.12
FBW (g)	24.88±1.97 <sup>c</sup>	28.58±0.76 <sup>b</sup>	29.58±0.72 <sup>b</sup>	26.88±1.94 <sup>c</sup>	31.76±1.09 <sup>a</sup>
WG (g)	20.43±2.18 <sup>d</sup>	24.05±0.76 <sup>b</sup>	25.05±0.72 <sup>b</sup>	22.40±1.88 <sup>c</sup>	27.36±1.02 <sup>a</sup>
ADG(g day <sup>-1</sup> )	0.34±0.03 <sup>d</sup>	0.40±0.01 <sup>bc</sup>	0.41±0.01 <sup>b</sup>	0.37±0.03 <sup>c</sup>	0.45±0.01 <sup>a</sup>
SGR (% day <sup>-1</sup> )	1.24± 0.09 <sup>c</sup>	1.33± 0.03 <sup>b</sup>	1.35±0.04 <sup>b</sup>	1.29±0.05 <sup>bc</sup>	1.43±0.02 <sup>a</sup>
FCR (%)	1.79±0.16 <sup>a</sup>	1.51±0.04 <sup>b</sup>	1.37±0.04 <sup>c</sup>	1.55±0.18 <sup>b</sup>	1.28±0.06 <sup>c</sup>
FL(cm)	11.60±0.43 <sup>b</sup>	11.98±0.31 <sup>b</sup>	12.23±0.61 <sup>ab</sup>	12.01±0.51 <sup>b</sup>	12.66±0.59 <sup>a</sup>
CF	1.76±0.20	1.87±0.13	1.84±0.15	1.82±0.28	1.79±0.15
SR (%)	96.65±4.73	96.65±4.73	96.65±4.73	91.65±2.33	91.65±2.33
HSI (%)	1.74±0.27 <sup>ab</sup>	1.67±0.17 <sup>b</sup>	1.97±0.19 <sup>a</sup>	2.02±0.22 <sup>a</sup>	2.22±0.25 <sup>a</sup>

Data are expressed as means±standard deviations. Values with the same superscripts of the same row are not significantly different ( $p>0.05$ ). Where, IBW= initial body weight, FBW=final body weight, WG=weight gain, ADG=average daily gain, SGR=specific growth rate, FCR=food conversion ratio, FL=fish length, CF=condition factor, HSI=Hepatosomatic Index.

**Table 3: Some biochemical parameters, liver and kidney functions of *Oreochromis niloticus* 60 days after dietary *Moringa oleifera* aqueous extract supplementation.**

parameters	Treatments				
	Dietary <i>M. oleifera</i> extract mg kg <sup>-1</sup> diet				
	0 (Control)	50	100	200	400
Total protein (g)	4.0±0.80 <sup>b</sup>	5.16±1.26 <sup>ab</sup>	5.15±1.05 <sup>ab</sup>	5.23±1.43 <sup>ab</sup>	6.20±1.89 <sup>a</sup>
Serum IgM (µg ml <sup>-1</sup> )	117.34±14.24 <sup>b</sup>	127.19±34.94 <sup>ab</sup>	125.56±16.16 <sup>ab</sup>	150.15±44.86 <sup>a</sup>	112.77±11.62 <sup>b</sup>
Albumin (g dl <sup>-1</sup> )	1.88±0.14	1.96± 0.52	1.89± 0.39	2.10±0.34	2.03±0.52
AST (U L <sup>-1</sup> )	58.08±12.64 <sup>a</sup>	56.08±14.45 <sup>a</sup>	56.50±11.00 <sup>a</sup>	55.72±9.98 <sup>a</sup>	44.00±6.28 <sup>b</sup>
ALT ( U L <sup>-1</sup> )	39.33±19.93 <sup>a</sup>	40.16± 14.62 <sup>a</sup>	38.26±16.13 <sup>a</sup>	40.98±18.64 <sup>a</sup>	31.33±10.35 <sup>b</sup>
Urea (mg dl <sup>-1</sup> )	13.14±1.78 <sup>a</sup>	10.89± 2.60 <sup>ab</sup>	9.10±1.57 <sup>b</sup>	11.16±3.37 <sup>ab</sup>	11.17±2.64 <sup>ab</sup>
Creatinine (mg dl <sup>-1</sup> )	0.73±0.09	0.73±0.10	0.71±0.08	0.65±0.20	0.80±0.11
Blood glucose (mg dl <sup>-1</sup> )	115.33±7.17 <sup>a</sup>	89.16±16.57 <sup>b</sup>	79.00±13.79 <sup>b</sup>	133.00±31.90 <sup>a</sup>	85.50±19.36 <sup>b</sup>

Data are expressed as means ± standard deviations. Values with the same superscripts in the same row are not significantly different ( $p>0.05$ ).

Additionally, the dietary MOAE received fish groups showed a pronounced decrease in the SOD, GPX, G6PDH and NO activities (Table 4).


The dietary MOAE supplementation had also triggered a substantial ( $p < 0.05$ ) deviation in the cortisol levels of the *M. oleifera* fed fish during the pre and post-hypoxic stress period (Fig. 1). Regarding the pre-hypoxic condition period; the MOAE supplemented fish (G1; 50 mg kg<sup>-1</sup> diet) exhibited a significantly lower cortisol level as matched to both the control fish group and G<sub>4</sub> (200 mg MOAE kg<sup>-1</sup> diet).

On the contrary to pre-hypoxic condition, elevated serum cortisol levels were equally assessed as a hypoxia-related response in all treated fish groups 30 minute's post-hypoxic stress. Nevertheless, the MOAE supplemented diets decreased the hypoxia-related stress in hypoxic tilapia as conveyed by the gradual drop in the cortisol level two hours after induction of hypoxia. The lowest serum cortisol values were detected in the G<sub>4</sub> and G<sub>5</sub>.

**Table 4: Antioxidant enzyme activities in *Oreochromis niloticus* 60 days after dietary *Moringa oleifera* aqueous extract supplementation.**

Parameters	Treatments				
	Dietary <i>M. oleifera</i> extract mg kg <sup>-1</sup>				
	0 (Control)	50	100	200	400
SOD (U mg <sup>-1</sup> )	16.69±0.82 <sup>a</sup>	12.16±2.63 <sup>b</sup>	7.24±2.02 <sup>c</sup>	7.14± 2.79 <sup>c</sup>	5.56±0.60 <sup>c</sup>
G6PD (mU mg <sup>-1</sup> )	6.21±3.27 <sup>a</sup>	6.10±1.05 <sup>a</sup>	2.05±1.08 <sup>b</sup>	1.56±0.52 <sup>b</sup>	2.17±1.32 <sup>b</sup>
GPX (mU mg <sup>-1</sup> )	4.56±0.56 <sup>a</sup>	4.11±0.26 <sup>a</sup>	2.29±0.28 <sup>b</sup>	2.11±0.28 <sup>b</sup>	1.13±0.11 <sup>c</sup>
(NO) activity (μ mol L <sup>-1</sup> )	26.43±3.55 <sup>a</sup>	18.97± 8.82 <sup>abc</sup>	20.07±5.25 <sup>ab</sup>	15.35±4.53 <sup>bc</sup>	9.01±0.96 <sup>c</sup>

Data are expressed as means ± standard deviations. Values with the same superscripts in the same row are not significantly different ( $p > 0.05$ ).



**Figure 1: Serum cortisol levels of *Oreochromis niloticus* challenged with acute hypoxia after 60 days of dietary *Moringa oleifera* aqueous extract supplementation. Values with the same superscript letters at initial, 30 ms and 2 hours post-hypoxic stage are not significantly different ( $p > 0.05$ ) between different treated groups.**



## Discussion

MOAE was recently verified to possess a valuable range of active substances. They possess growth promoting, antioxidant, tissue protective (liver, kidneys, heart and testes), anti-stressor, and immunomodulatory activities in fish (Alishahi *et al.*, 2010; Stadlander *et al.*, 2013). This study clarified that feeding tilapia with MOAE containing diet enhanced the growth performance of *O. niloticus* fingerlings distinctly. The FCR was improved as the dietary inclusion level of MOAE increased. Fish received 400 mg MOAE kg<sup>-1</sup> had a markedly high WG, SGR, and ADG. The ethanolic extract of *M. oleifera* flower (250 and 500 mg kg<sup>-1</sup> diet) as well had improved tilapia growth (Tekle and Sahu, 2015). Also, a growth-promoting potential of *M. oleifera* seed protein extracts was reported by Stadlander *et al.* (2013) when added to tilapia diet at 400 mg kg<sup>-1</sup> in an eight weeks experimental period. Not only this but also the dried powder of *M. oleifera* leaves (10 mg kg diet<sup>-1</sup>) was able to promote the growth of juveniles of *Penaeus indicus* as reported by Rayes (2013).

*M. oleifera* leaves is said to contain vast amount of immuno- nutritional components such as proteins, lipids, vitamins, enzymes, minerals, sugar, saponin and salicylates and other active constituents (Leone *et al.*, 2015). Moreover, Heidarieh *et al.* (2013) indicated that *M. oleifera* had enhanced the gastrointestinal morphology of *O. mykiss* (increased villi length) which improved the food digestibility and absorption capacity. Herein, the

increment in growth and diet utilization could be related to its immuno- nutritional components and the high feed digestibility, absorption and assimilation ability, through the augmented digestive enzymes and healthy intestinal microflora boosted by the *M. oleifera* prebiotic activity.

In the meantime, the usage of either raw *M. oleifera* or its extracts as a dietary supplement or protein replacement has shown variable results. Feeding Moringa leaf meal to Nile tilapia at a dietary inclusion level up to 8-10% of the diets hindered tilapia growth (Richter *et al.*, 2003; Yuangsoi and Charoenwattanasak, 2011; Abo-State *et al.*, 2014). Similarly, Afuang *et al.* (2003) incorporated methanol-extracted leaf meal containing fewer saponins and phenolics in the diets (at a 30% inclusion level) of tilapia fingerlings. The fish performance was not hindered, but the body protein content reduced. Whereas, the aqueous extracted leaf meal (15% of the diet) had reduced the feed intake, feed utilization and fish performance (Madalla, 2008). Also, Dongmeza *et al.* (2006) reported a considerable decrease in the Nile tilapia feed intake and consequently its growth after receiving different dietary moringa leaf extracts (tannin-reduced, saponin-reduced and saponin-enriched).

Measurement of the plasma biochemical parameters is a fundamental step to judge the health integrity of any aquatic species (Ferreira *et al.*, 2007). Plasma proteins carry out many functions, such as adjusting the water balance in fish

(Rudneva and Kovyrshina, 2011) and the protective effects implemented by the body for the elimination of any microorganisms (Gerwick *et al.*, 2002). This study displayed an increased level of plasma protein in the dietary MOAE supplemented fish. Similar conclusions were revealed by Alishahi *et al.* (2010) and Haghighi *et al.* (2014) who informed that 0.5% and 1% *M. oleifera* dietary supplementation had significantly improved the serum total protein and globulin in *Cyprinus carpio* and *O. mykiss* respectively. Conversely, the serum total protein levels obtained in this work could not agree with either Dotta *et al.* (2014) for *O. niloticus* (0.5% *M. oleifera*) nor Kavitha *et al.* (2012) for *Cyprinus carpio* (exposed to 124.0 mg L<sup>-1</sup> *M. oleifera* extract). They reported that *M. oleifera* dietary supplementation was unable to considerably elevate the total serum protein in those fish as an outcome of the metabolic degradation and utilization of protein.

Transaminase enzymes are of dynamic importance in protein and carbohydrate metabolism. Their levels (i.e. transaminases; AST and ALT) are used for judging the physiological and healthy state of aquatic species (Vutukuru *et al.*, 2007). Therefore, they are indicative of organ dysfunction, especially for hepatic dysfunction with a subsequent leakage of enzymes into the blood (Gabriel and George, 2005). The aminotransferase activities were comparable among all the treated groups except for the group fish supplemented with 400 mg MOAE kg diet<sup>-1</sup> which displayed a reasonable

diminishing in the activity aminotransferase enzymes. This result supports the previous studies that convey the potentiality of *M. oleifera* leaves extract to protect the membrane integrity of tilapia hepatocytes against stressors (Tekle and Sahu, 2015). Also, the increased growth rate conveyed in the current study give further support for the liver integrity as a vital organ implemented in the normal metabolic function of any aquatic organism.

Creatinine and urea are among the dominant biochemical indices particularly determined to assess renal condition (Gross *et al.*, 2005; Adeyemi and Akanji, 2012). The fish received 100 mg MOAE kg<sup>-1</sup> exhibited a significant decrease in the urea level. Such reduction may offer a further support of the nephron-protection activity of *M. oleifera* as reported by Anwar *et al.* (2007) and Sharma and Paliwal (2012). Also, the comparable creatinine levels distinguished in differently treated fish groups provide further support to the nephron-protective potentiality of MOAE. Nevertheless, our results are dissimilar to Mazumder *et al.* (1999) who clarified that a weekly dose (>46 mg kg<sup>-1</sup> b.wt extract) of *M. oleifera* methanol root extracts had impaired function of mice kidney. Oyagbemi *et al.* (2013) as well reported elevated serum creatinine levels in rats administered *M. oleifera* (200 and 400 mg kg<sup>-1</sup> b.w.).

Regarding the blood glucose levels verified in the current study; the MOAE supplemented diet had hypoglycemic effect in all of the treated fish groups. Similarly, fish received herbal additives

showed hypoglycemia (Metwally, 2009; Banaee *et al.*, 2011; Ojha *et al.*, 2014). MOAE had successfully governed the glucose level in rabbits (Makonnen *et al.*, 1997) and rats (Amin *et al.*, 2016). The moringa leaf juices contain  $\alpha$ -glucosidase and amylase inhibitors. These enzymes prevent the assimilation of glucose into absorbable metabolites, hence preventing the increase in blood glucose (Abdulkarim *et al.*, 2005).

On the contrary; diabetes inducing property was verified to green tea extract indicated by the observed hyperglycemia in green tea extract exposed Nile tilapia (Abdel-Tawwab *et al.*, 2010).

SOD and GPX are sensitive oxidative stress biomarkers. They are considered the core line of the body antioxidant defensive mechanism (Jiang *et al.*, 2009). SOD is a dynamic enzyme responsible for both scavenging ROS and protection of cells from being damaged by free radicals (Chien *et al.*, 2003). In our study, the MOAE supplementation considerably diminished the SOD and GPX activity, indicating that *M. oleifera* might have the capability to reduce peroxide radicals and converting it into oxygen and water due to an antioxidant potential (Atli and Canli, 2007). *M. oleifera* could also preserve the cell redox state by restricting the damaging effects of ROS (Halliwell and Gutteridge, 2007). Furthermore, the MOAE dietary supplementation resulted in reduced activity of other antioxidant enzymes. *M. oleifera* contains active biological compounds

(polyphenols, glycosides, anthocyanin, tannins and thiocarbamates). These active compounds expel free radicals, active antioxidant enzymes and inhibit oxidases (Luqman *et al.*, 2011) which give a further clarification for the *M. oleifera* associated antioxidant activity.

Furthermore, this study revealed that acutely hypoxic tilapia had exhibited an elevation in the cortisol level 30 minute's post-hypoxic stress. The elevated cortisol level was reduced distinctly two hours post-hypoxic stress in those fish received 200 and 400 mg MOAE kg<sup>-1</sup> feed. The retrieval towards normal cortisol level following MOAE pretreatment proposed hypocortisolemic effects of MOAE. Additionally, the potent antioxidants of Moringa supplemented diets was said to be useful in compromising the adversative properties of stress related hypoxia. Production of oxidative radicals (especially ROS) is thought to be boosted under hypoxic stress (Chandel and Budinger, 2007). Accordingly, these inclusion levels were possibly capable of blocking the hypercortisolemia elicited by the acute stress caused by hypoxia or efficiently restored the homeostatic equilibrium of the life helping physiological systems (Van Rijn and Reina, 2010). Hamed *et al.* (2015) also reported that extracts of Moringa leaves infiltrate to the cell membrane lipid bilayer, resulting in improved permeability, and ROS elimination. Hence these fish groups could successfully tolerate and recover the hypoxic stress than the others. This study verified the potentiality of *M. oleifera* aqueous extract as a growth

promoter, antistress and antioxidant for *O. niloticus* fingerlings. *M. oleifera* aqueous extracts are proposed to replace synthetic antimicrobials and growth promoters. This study is valuable to validate the safe administration of *M. oleifera* in commercial tilapia culture. In this concern, future studies are required to comprehend the influence of this plant extract in relieving chronic stress and its possible toxic effect as well. Feasibility study for its commercial usage is required too.

#### Acknowledgements

This work was supported by the department of Animal Husbandry and Animal Wealth Development, Faculty of Veterinary Medicine, Alexandria University, Egypt.

#### References

- Abdel-Tawwab, M., Ahmad, M.H., Seden, M.E. and Sakr, S.F., 2010.** Use of green tea, *Camellia sinensis* L., in practical diet for growth and protection of Nile tilapia, *Oreochromis niloticus* (L.), against *Aeromonas hydrophila* infection. *Journal of the World Aquaculture Society*, 41, 203-213.
- Abdulkarim, S.M., Long, K., Lai, O.M., Muhammad, S.K.S. and Ghazali, H.M., 2005.** Some physico-chemical properties of *Moringa oleifera* seed oil extracted using solvent and aqueous enzymatic methods. *Food Chemistry*, 93(2), 253-263.
- Abo-State, H., Hammouda, Y., El-Nadi, A. and AboZaid, H., 2014.** Evaluation of feeding raw moringa (*Moringa oleifera* Lam.) leaves meal in Nile tilapia fingerlings (*Oreochromis niloticus*) diets. *Global Veterinaria*, 13(1), 105-111.
- Adeyemi, O.S. and Akanji, M.A., 2012.** *Psidium guajava* leaf extract: Effects on rat serum homeostasis and tissue morphology. *Comparative Clinical Pathology*, 21(4), 401-407.
- Afuang, W., Siddhuraju, P. and Becker, K., 2003.** Comparative nutritional evaluation of raw methanol extracted residues and methanol extracts of moringa (*Moringa oleifera* Lam.) leaves on growth performance and feed utilization in Nile tilapia (*Oreochromis niloticus* L.). *Aquaculture Research*, 34(13), 1147-1159.
- Alishahi M, Ranjbar, M., Ghorbanpoor, M., Peyghan, R. and Mesbah, M., 2010.** Effects of dietary *Aloe vera* on some specific and nonspecific immunity in the common carp (*Cyprinus carpio*). *International Journal of Veterinary Research*, 4(3), 189-195.
- Amin, A.Y., El Tobgy, K.M.K., Salam, A. and Hemat, S., 2016.** Phytochemical detection and Therapeutical properties of *Moringa oleifera* leaves. *International Journal of Chem Tech Research*, 9(9), 156-168.
- Anwar, F., Latif, S., Ashraf, M. and Gilani, A.H., 2007.** *Moringa oleifera*: A food plant with multiple medicinal uses. *Phytotherapy Research*, 21(1), 17-25.

- Atli, G. and Canli, M., 2007.** Enzymatic responses to metal exposures in a freshwater fish *Oreochromis niloticus*. *Comparative Biochemistry and Physiology Part C: Toxicology and Pharmacology*, 145(2), 282-287.
- Banaee, M., Sureda, A., Mirvaghefi, A.R. and Rafei, G.R., 2011.** Effects of long-term silymarin oral supplementation on the blood biochemical profile of rainbow trout (*Oncorhynchus mykiss*). *Fish Physiology and Biochemistry*, 37(4), 885-896.
- Bautista, J., Garrido-Pertierra, A. and Soler, G., 1988.** Glucose-6-phosphate dehydrogenase from *Dicentrarchus labrax* liver: Kinetic mechanism and kinetics of NADPH inhibition. *Biochimica et Biophysica Acta (BBA)-General Subjects*, 967(3), 354-363.
- Bulfon, C., Volpatti, D. and Galeotti, M., 2015.** Current research on the use of plant-derived products in farmed fish. *Aquaculture Research*, 46(3), 513-551.
- Burnett, B.P., Pillai, L., Bitto, A., Squadrito, F. and Levy, R.M., 2011.** Evaluation of CYP450 inhibitory effects and steady-state pharmacokinetics of genistein in combination with cholecalciferol and citrated zinc bisglycinate in postmenopausal women. *International Journal of Women's Health*, 3(1), 139-150.
- Chakraborty, S.B. and Hancz, C., 2011.** Application of phytochemicals as immunostimulant, antipathogenic and antistress agents in finfish culture. *Reviews in Aquaculture*, 3(3), 103-119.
- Chandel, N.S. and Budinger, G.S., 2007.** The cellular basis for diverse responses to oxygen. *Free Radical Biology and Medicine*, 42(2), 165-174.
- Chien, Y.H., Pan, C.H. and Hunter, B., 2003.** The resistance to physical stresses by *Penaeus monodon* juveniles fed diets supplemented with astaxanthin. *Aquaculture*, 216(1), 177-191.
- Cooper, G.R. and McDaniel, V., 1970.** The determination of glucose by the ortho-toluidine method (filtrate and direct procedure). *Standard Methods of Clinical Chemistry*. 6, 159-170.
- Dongmeza, E., Siddhuraju, P., Francis, G. and Becker, K., 2006.** Effects of dehydrated methanol extracts of moringa (*Moringa oleifera* Lam.) leaves and three of its fractions on growth performance and feed nutrient assimilation in Nile tilapia (*Oreochromis niloticus* (L.)). *Aquaculture*, 261(1), 407-422.
- Dotta, G., de Andrade, J.I.A., Gonçalves, E.L.T., Brum, A., Mattos, J.J., Maraschin, M. and Martins, M.L., 2014.** Leukocyte phagocytosis and lysozyme activity in Nile tilapia fed supplemented diet with natural extracts of propolis and *Aloe barbadensis*. *Fish and Shellfish Immunology*, 39(2), 280-284.
- Drupt, F., Paris, M., Frydman, A. and Leclerc, M., 1974.** Serum albumin assay by bromocresol green method: Application to different automatic apparatus. *Annales*

- Pharmaceutiques Francaises*, (Vol. 32(5), 249).
- FAO., 2014.** Food and agricultural organisation of the United Nations. The state of world Fisheries and Aquaculture 2014. Rome. Italy.
- Fawcett, J.K. and Scott, J.E., 1960.** Enzymatic, colorimetric method for determination urea in serum, plasma and urine. *Journal of Clinical Pathology*, 13, 156–162.
- Fernandez, T.J., 1990.** Medicinal plants for Haemophilus gallinarum infection in chickens. *ASEAN Journal on Science and Technology for Development*, 7(22), 99-107.
- Ferreira, J.G., Hawkins, A.J.S. and Bricker, S.B., 2007.** Management of productivity, environmental effects and profitability of shellfish aquaculture—the Farm Aquaculture Resource Management (FARM) model. *Aquaculture*, 264(1), 160-174.
- Gabriel, U.U. and George A.O.I., 2005.** Plasma enzymes in *Clarias gariepinus* exposed to chronic levels of roundup (glyphosate). *Journal of Ecology and Environment*, 23(2), 271–276.
- Gerwick, L., Steinhauer, R., Lapatra, S., Sandell, T., Ortuno, J., Hajiseyedjavadi, N. and Bayne, C.J., 2002.** The acute phase response of rainbow trout (*Oncorhynchus mykiss*) plasma proteins to viral, bacterial and fungal inflammatory agents. *Fish and Shellfish Immunology*, 12(3), 229-242.
- Gross, J.L., De Azevedo, M.J., Silveiro, S.P., Canani, L.H., Caramori, M.L. and Zelmanovitz, T., 2005.** Diabetic nephropathy: Diagnosis, prevention, and treatment. *Diabetes Care*, 28(1), 164-176.
- Haghighi, M., Rohani, M.S., Pourmoghim, H., Toliat, T., Samadi, M., Tavoli, M., Islami, M. and Yusefi, R., 2014.** Haemato-immunological indices in rainbow trout (*Oncorhynchus mykiss*) fry fed with *Aloe vera* extract supplemented feed. *Journal of Coastal Life Medicine*, 2, 350-356.
- Halliwell, B. and Gutteridge, J.M.C., 2007.** Antioxidant defences: Endogenous and diet derived. *Free Radicals in Biology and Medicine*, 4, 79-186.
- Hammed, A.M ,Amosu A.O, Awe, A.F., and Gbadamosi, F.F., 2015.** Effects of *Moringa oleifera* leaf extracts on bacteria (*Aeromonas hydrophila*) infected adults African mud cat fish. *International Journal of Current Research*, 7(10), 22117-22122.
- Heidarieh, M., Mirvaghefi, A.R., Sepahi, A., Sheikhzadeh, N., Shahbazfar, A.A. and Akbari, M., 2013.** Effects of dietary *Aloe vera* on growth performance, skin and gastro-intestine morphology in rainbow trout (*Oncorhynchus mykiss*). *Turkish Journal of Fisheries and Aquatic Sciences*, 13, 367-373.
- Husdan, H. and Rapoport, A., 1968.** Estimation of creatinine by the Jaffe reaction. *Clinical Chemistry*, 14(3), 222-238.
- Huyghebaert, G., Ducatelle, R. and Van Immerseel, F., 2011.** An

- update on alternatives to antimicrobial growth promoters for broilers. *The Veterinary Journal*, 187(2), 182-188.
- Jiang, W.D., Feng, L., Liu, Y., Jiang, J. and Zhou, X.Q., 2009.** Myo-inositol prevents oxidative damage, inhibits oxygen radical generation and increases antioxidant enzyme activities of juvenile Jian carp (*Cyprinus carpio* var. Jian). *Aquaculture Research*, 40(15), 1770-1776.
- Kautsky, N., Rönnbäck, P., Tedengren, M. and Troell, M., 2000.** Ecosystem perspectives on management of disease in shrimp pond farming. *Aquaculture*, 191(1), 145-161.
- Kavitha, C., Ramesh, M., Kumaran, S.S. and Lakshmi, S.A., 2012.** Toxicity of *Moringa oleifera* seed extract on some hematological and biochemical profiles in a freshwater fish, *Cyprinus carpio*. *Experimental and Toxicologic Pathology*, 64(7), 681-687.
- Knobil, K., 1980.** Hormonal researches. *Recent Progress in Hormone Research*, 36, 52-58.
- Leone, A., Spada, A., Battezzati, A., Schiraldi, A., Aristil, J. and Bertoli, S., 2015.** Cultivation, genetic, ethnopharmacology, phytochemistry and pharmacology of *Moringa oleifera* leaves: An overview. *International Journal of Molecular Sciences*, 16(6), 12791-12835.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J., 1951.** Protein measurement with the Folin phenol reagent. *Journal of Biological Chemistry*, 193(1), 265-275.
- Luqman, S., Srivastava, S., Kumar, R., Maurya, A.K. and Chanda, D., 2011.** Experimental assessment of *Moringa oleifera* leaf and fruit for its antistress, antioxidant, and scavenging potential using in vitro and in vivo assays. *Evidence-Based Complementary and Alternative Medicine*, 2012, 1-12.
- Madalla, N., 2008.** Novel feed ingredients for Nile tilapia (*Oreochromis niloticus* L.). In: PhD Thesis, Inst. Aquaculture, Univ. Stirling, Scotland, UK.
- Makonnen, E., Hunde, A. and Damecha, G., 1997.** Hypoglycaemic effect of *Moringa stenopetala* aqueous extract in rabbits. *Phytotherapy Research*, 11(2), 147-148.
- Mazumder, U.K., Gupta, M., Chakrabarti, S. and Pal, D., 1999.** Evaluation of hematological and hepatorenal functions of methanolic extract of *Moringa oleifera* Lam. root treated mice. *Indian Journal of Experimental Biology*, 7, 21-25.
- Mbikay, M., 2012.** Therapeutic potential of *Moringa oleifera* leaves in chronic hyperglycemia and dyslipidemia: A review. *Frontiers in Pharmacology*, 3, 24.
- Menz, J., Schneider, M. and Kümmerer, K., 2015.** Usage pattern-based exposure screening as a simple tool for the regional priority-setting in environmental risk assessment of veterinary antibiotics:

- A case study of north-western Germany. *Chemosphere*, 12, 42-48.
- Metwally, M.A.A., 2009.** Effects of garlic (*Allium sativum*) on some antioxidant activities in tilapia nilotica (*Oreochromis niloticus*). *World Journal of Fish and Marine Sciences*, 1(1), 56-64.
- Moss, S.M., Moss, D.R., Arce, S.M., Lightner, D.V. and Lotz, J.M., 2012.** The role of selective breeding and biosecurity in the prevention of disease in penaeid shrimp aquaculture. *Journal of Invertebrate Pathology*, 110(2), 247-250.
- Nikkon, F., Saud, Z.A., Rehman, M.H. and Haque, M.E., 2003.** In vitro antimicrobial activity of the compound isolated from chloroform extract of *Moringa oleifera* Lam. *Pakistan Journal of Biological Science*, 22, 1888-1890.
- Nishikimi, M., Rao, N.A. and Yagi, K., 1972.** The occurrence of superoxide anion in the reaction of reduced phenazine methosulfate and molecular oxygen. *Biochemical and Biophysical Research Communications*, 46(2), 849-854.
- Ojha, M.L., Chadha, N.K., Saini, V.P., Damroy, S., Gupta, C.P. and Savant, P.B., 2014.** Effect of ethanolic extract of *Pedaliium murex* on growth and haemato-immunological parameters of *Labeo rohita*. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences*, 84(4), 997-1003.
- Oyagbemi, A.A., Omobowale, T.O., Azeez, I.O., Abiola, J.O., Adedokun, R.A. and Nottidge, H.O., 2013.** Toxicological evaluations of methanolic extract of *Moringa oleifera* leaves in liver and kidney of male Wistar rats. *Journal of Basic and Clinical Physiology and Pharmacology*, 24(4), 307-312.
- Paglia, D.E. and Valentine, W.N., 1967.** Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *Journal of Laboratory and Clinical Medicine*, 7, 158-169.
- Rayes, A.H., 2013.** Study on the effect of dietary probiotic bacteria *Arthrobacter* species,  $\beta$ -1, 3 glucan and *Moringa oleifera* leaf on protection of *Penaeus indicus* Juveniles from pathogenic *Vibrio harveyi*. *Researcher*, 5(1), 24-31.
- Reitman, S. and Frankel, S., 1957.** A colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases. *American Journal of Clinical Pathology*, 28(1), 56-63
- Reverter, M., Bontemps, N., Lecchini, D., Banaigs, B. and Sasal, P., 2014.** Use of plant extracts in fish aquaculture as an alternative to chemotherapy: Current status and future perspectives. *Aquaculture*, 433, 50-61.
- Richter, N., Siddhuraju, P. and Becker, K., 2003.** Evaluation of nutritional quality of moringa (*Moringa oleifera* Lam.) leaves as an alternative protein source for Nile tilapia (*Oreochromis niloticus* L.). *Aquaculture*, 217(1), 599-611.
- Rudneva, I.I. and Kovyreshina, T.B., 2011.** Comparative study of electrophoretic characteristics of



- serum albumin of round goby *Neogobius melanostomus* from Black Sea and Azov Sea. *International Journal of Science and Nature*, 1, 131-136.
- SAS., 2008.** Statistical analysis system. User's Guide. SAS INT., Cary, NC. USA.
- Schreck, C.B., 2000.** Accumulation and long-term effects of stress in fish. In: Moberg, G.P., Mench, J.A. (Eds.), *The Biology of Animal Stress: Basic Principles and Implications for Animal Welfare*. CABI Publishing, CAB International, Oxon and New York. pp. 147–158.
- Sharma, V. and Paliwal, R., 2012.** Chemo protective role of *Moringa oleifera* and its isolated saponin against DMBA induced tissue damage in male mice: A histopathological analysis. *International Journal of Drug Development and Research*, 4(4), 215-228.
- Simi, S., Peter, V.S. and Peter, M.S., 2016.** Zymosan-induced immune challenge modifies the stress response of hypoxic air-breathing fish (*Anabas testudineus* Bloch): Evidence for reversed patterns of cortisol and thyroid hormone interaction, differential ion transporter functions and non-specific immune response. *General and Comparative Endocrinology*. 251, 94-108. <http://doi.org/10.1016/j.ygcen.2016.11.009>
- Stadtlander, T., Sander, C., Kumar, V., Makkar, H. and Becker, K., 2013.** Effects of *Moringa oleifera* Lam. dietary seed protein extracts on growth, nutrient utilization and blood parameters in common carp (*Cyprinus carpio*, L.) and Nile tilapia (*Oreochromis niloticus*, L.). *Planta Medica*, 79(13), PL21.
- Suarez, M., Entenza, J.M., Doerries, C., Meyer, E., Bourquin, L., Sutherland, J., Marison, I., Moreillon, P. and Mermod, N., 2003.** Expression of a plant-derived peptide harboring water-cleaning and antimicrobial activities. *Biotechnology and Bioengineering*, 81(1), 13-20.
- Tekle, E.W. and Sahu, N.P., 2015.** Growth and immunodulatory response of Nile tilapia *Oreochromis niloticus* fingerlings to ethanolic extract of *Moringa oleifera* flower. *International Journal of Scientific and Research Publications*, 5(7), 285-296.
- Van Bezooijen, R.L., Que, I., Ederveen, A.G., Kloosterboer, H.J., Papapoulos, S.E. and Lowik, C.W., 1998.** Plasma nitrate+ nitrite levels are regulated by ovarian steroids but do not correlate with trabecular bone mineral density in rats. *Journal of Endocrinology*, 159(1), 27-34.
- Van Rijn, J.A. and Reina, R.D., 2010.** Distribution of leukocytes as indicators of stress in the Australian swellshark, *Cephaloscyllium laticeps*. *Fish and Shellfish Immunology*, 29(3), 534-538.
- Vutukuru, S.S., Pauleena, J.S., Rao, J.V. and Anjaneyulu, Y., 2007.** Architectural changes in the gill

morphology of the freshwater fish, *Esomus danricus* as potential biomarkers of copper toxicity using automated video tracking system. *Environmental Bioindicators*, 2(1), 3-14.

**Wang, T., Lefevre, S., Van Cong, N. and Bayley, M., 2009.** The effects of hypoxia on growth and digestion. *Fish Physiology*, 27, 361-396.

**Yuangsoi, B. and Charoenwattanasak, S., 2011.** Utilization of moringa (*Moringa oleifera* Lam.) leaf on growth performance and protein digestibility in Tilapia (*Oreochromis niloticus* L.). Proceedings of the 49th Kasetsart University Annual Conference, Kasetsart University, Thailand, 1-4 February, 2011. Vol. 3. Subject: Fisheries Kasetsart University. pp. 317-326