



Official Journal Issued by
Faculty of
Veterinary Medicine

Benha Veterinary Medical Journal

Journal homepage: <https://bvmj.journals.ekb.eg/>



Since 1990

Original Paper

Assessment of histamine residues in smoked and salted fish

Mohamed A. Hassan¹, Fahim A. Shaltout², Nabila E. El-sheikh³, Naglaa M. Sakr⁴

^{1,2} Food Hygiene Department, Faculty of Veterinary Medicine, Benha University, Egypt.

^{3,4} Animal Health Research Institute, Tanta, Egypt

ARTICLE INFO

Keywords

Histamine
Salted sardine
Smoked fish

ABSTRACT

Histamine is a member of a group of compounds known as biogenic amines normally produced by decarboxylation of free amino acids and are present in a variety of foods. In the present study a total of 90 random samples of salted and smoked fish products represented by fesiekh, salted sardine, and smoked herring (30 of each) were collected from different fish markets in Gharbia governorate, Egypt and examined for the presence of histamine by ELISA. The results recorded that the histamine mean values in examined fish samples were 20.76 ± 0.54 , 15.49 ± 0.31 and 9.82 ± 0.26 mg/kg. Unaccepted samples were 53.3%, 36.7% and 30% for fesiekh, sardine and smoked herring, respectively.

1. INTRODUCTION

Fish is a very important source of protein especially in Egypt. Fish and fish products are one of the most important food stuffs as they are one of the cheapest sources of animal protein. Fish are enriched with essential minerals, vitamins, and unsaturated fatty acids (El-Moselhy, 2000). Histamine is known as a biogenic amine which is low molecular weight and possesses biological activity (Tombelli and Mascini, 1998). The levels of histamine have been suggested as rapid fish spoilage indicators (Tombelli and Mascini, 1998; Patange et al., 2005). Gram-negative histamine producing bacteria are more common in fish. A wide range of Gram-negative bacteria can produce histamine in fish, but the major types are mesophilic enteric and marine bacteria. *Morganella morganii*, *Morganella psychrotolerans*, *Photobacterium damsela*, *Photobacterium phosphoreum*, *Raoultella planticola*, *Hafnia alvei* were reported as histamine formers. In the case of fermented sea food, *Staphylococcus* spp. and *Tetragenococcus* spp. (Surya et al., 2019). These types of bacteria naturally present on the gills, external surfaces and in the gut of live saltwater fish with no harm to the fish. Up on death, the defense mechanism of the fish no longer inhibits bacterial growth in the muscle tissue and histamine forming bacteria may start to grow resulting in the formation of biogenic amines (FDA, 2011). Scombroid poisoning is a form of toxicity caused by the ingestion of spoiled dark-flesh fishes, mainly of the scombroid family. The clinical picture is secondary to histamine toxicity, manifested as flushing, headache, palpitations, and abdominal cramps (Ferran and Yébenes, 2006). Inadequate cooling following harvest promotes bacterial histamine production and can result in outbreaks of scombroid poisoning (Hungerford, 2010) which results from the ingestion of histamine-contaminated fish of the scombroid fish including tuna, mackerel, and non-

scombroid fish include sardine, herring and anchovy (Feng et al., 2016). The symptoms of scombroid poisoning appear within a few minutes after eating fish of scombridae family and related species. The first symptoms are cutaneous, with flush, pruritus, and erythema of the face and trunk having an urticarial appearance, together with faintness. Gastrointestinal symptoms include nausea, vomiting, abdominal cramps and occasionally diarrhea (Harmelin et al., 2018).

This work aimed to determine histamine residue in salted and smoked fish collected from different fish markets in Gharbia governorate, Egypt by using ELISA technique to assess quality of fish.

2. MATERIAL AND METHODS

2.1. Collection of samples:

A total of 90 random samples of salted and smoked fish products represented by fesiekh, salted sardine, and smoked herring (30 of each) were collected from different fish markets in Gharbia governorate, Egypt. The collected samples were labeled and preserved individually in an insulated ice box as well as transferred to the laboratory as quickly as possible.

2.2. Determination of histamine by ELISA:

2.2.1. Sample preparation and acylation:

Pipette 25 μ l of standards, 25 μ l of controls, 25 μ l of plasma samples, 10 μ l of fish samples, or 50 μ l of supernatant from the release test into the respective wells of the Reaction Plate. 25 μ l of Acylation Buffer were added to all wells. 25 μ l of Acylation Reagent were added to all wells. Incubated for 45 min at RT (20-25 °C) on a shaker (approx. 600 rpm). 200 μ l of distilled water were added to all wells. Incubated for 15 min. at RT (20-25°C) on a shaker (approx. 600 rpm). 25 μ l of the prepared standards, controls, and samples were

* Corresponding author: Naglaa Mahmoud Sakr. Animal Health Research Institute, Tanta, Egypt
E.mail: Naglaa.sakr333@gmail.com

taken for the Histamine ELISA. For the release test the Histamine Release supplementary kit (available for purchase separately, cat. no. BA E-1100) must be used.

2.2.2. Histamine ELISA:

25 µl of the acylated standards, controls, and samples were pipetted into the appropriate wells of the Histamine Microtiter Strips. 100 µl of the Histamine Antiserum were pipetted into all wells and cover plate with adhesive foil. Incubated for 3 hours at RT (20-25°C) on a shaker (approx. 600 rpm).

Alternatively: shake the Histamine Micro titer Strips briefly by hand and incubate for 15-20 hrs at 2-8°C.

The foil was removed. The contents of the wells were discarded or aspirated, and each well was washed 4 times thoroughly with 300 µl Wash Buffer. Blotted dry by tapping the inverted plate on absorbent material.

100 µl of the Enzyme Conjugate was pipetted into all wells. Incubated for 30 min at RT (20-25 °C) on a shaker (approx. 600 rpm).

The contents of well were discarded or aspirated, and each well was washed 4 times thoroughly with 300 µl. Wash Buffer. Blotted dry by tapping the inverted plate on absorbent material.

100 µl of the Substrate were pipetted into all wells and incubate for 20-30 min at RT (20-25 °C) on a shaker (approx. 600 rpm). Avoid exposure to direct sunlight.

100 µL of the Stop Solution were pipetted to each well and shake the microtiter plate to ensure a homogeneous distribution of the solution.

The absorbance of the solution in the wells was read within 10 minutes, using a microplate reader set to 450 nm with a reference wavelength between 620 nm and 650 nm.

3. Statistical Analysis

Analysis of Variance (ANOVA) test was applied for statistical evaluation of the obtained results of each detected residue in the examined samples of salted and smoked fish products according to Feldman et al. (2003).

3. RESULTS

It is evident from the results recorded in table (1) that the Histamine values in examined fish samples were varied from 2.9-36.1 mg/kg with an average of 20.76 ± 0.54 for Fesiekh; 2.2-29.8 mg/kg with an average of 15.49 ± 0.31 for sardine and 1.4-21.7 mg/kg with an average of 9.82 ± 0.26 for smoked herring. The differences between the examined samples of different fish species were high significant ($P < 0.01$). Table (2) revealed that 46.7%, 73.3% and 70% of the examined samples of Fesiekh, sardine and smoked herring were accepted. However, 53.3%, 36.7% and 30% of such samples were unaccepted, respectively.

Table 1 Concentrations of histamine (mg/Kg) in the examined samples of salted and smoked fish (n=30).

Fish products	Min	Max	Mean \pm S.E*
Fesiekh	2.9	36.1	20.76 ± 0.54
Sardine	2.2	29.8	15.49 ± 0.31
Smoked herring	1.4	21.7	9.82 ± 0.26

S.E* = Standard error of mean.

Table 2 Validity of the examined salted and smoked fish based on their histamine contents (n=30).

Fish products	MRL (mg/Kg) *	Accepted samples		Unaccepted samples	
		No.	%	No.	%
Fesiekh	20	14	46.7	16	53.3
Sardine	20	19	73.3	11	36.7
Smoked herring	20	21	70	9	30
Total (90)		54	60	36	40

* Egyptian Organization for Standardization "EOS" (2010).

4. DISCUSSION

The results recorded that the histamine mean values in examined fish samples were 20.76 ± 0.54 , 15.49 ± 0.31 and 9.82 ± 0.26 mg/kg, unaccepted samples were 53.3%, 36.7% and 30% for fesiekh, sardine and smoked herring, respectively.

Results of histamine in fesiekh nearly similar to those obtained by (Edris et al., 2014) (18.06 ± 0.99 mg/kg) who investigated histamine in 90 samples of fesiekh, sardine and melloha (30 of each) collected from different retail markets. Lower than that recorded by Nader et al. (2016) (33.21 ± 1.15 mg/100g), who investigated histamine in 20 samples of fesiekh were collected from markets in kafr El- Sheikh. Higher than that recorded by Elshafey-Wesam et al. (2018) (16.80 ± 0.31 mg/kg). measured histamine in 15 samples of Mugil cephalus.

Results of histamine in sardine lower than (Walaa, 2016) (29.65 ± 1.41 ppm) estimated the level of histamine in sardine samples using HPLC, Edris et al. (2014) (23.51 ± 1.21 mg/kg) who investigated histamine in 30 samples of sardine collected from different retail markets, Nader et al. (2016) (28.14 ± 1.02 mg/100g) and Elshafey-Wesam et al. (2018) (28.74 ± 0.52 mg/kg).

Results of histamine in smoked herring lower than Nader et al. (2016) (23.12 ± 0.86 mg /100g).

Fish commonly implicated in histamine fish poisoning include both scombroid (mackerel, tuna and saury) and non-scombroid (sardine, anchovies, blue fish, as they contain large amount of free histadine (Lehane and Olley, 2000). Sardines characterized by the presence of high levels of free histamine in their muscle, also according to the season of the year, genetics, environment, food, sex, physiological stage, storage period and sampled tissue (Lee et al., 2012). High levels of biogenic amines could be prevented by the application of good hygiene practices and proper temperature during handling, delivery and storage (Visciano et al., 2012). Biogenic amines could be used as quality index, once they are formed by bacterial activity and are resistant to thermal treatment, thus reflecting the quality of the raw material and hygienic conditions of food processing (Sagratiini et al., 2012). Acceptability of examined fish samples based on their levels of histamine according to EOS (2010) the results revealed that 46.7%, 73.3% and 70% of the examined samples of fesiekh, sardine and smoked herring were accepted, however, 53.3%, 36.7% and 30% of such samples were unaccepted, respectively. Unfortunately, unhygienic practices, insufficient refrigeration cause increase susceptible to contamination by BAs-producing microorganisms and other spoilage bacteria (proteolytic and lipolytic) leading to rapid spoilage and outbreaks of fish poisoning.

5. CONCLUSION

As conclusion, histamine is the main marker for the evaluation of quality and safety of fish. In addition, the application of good hygiene practices and proper temperatures during handling, delivery and storage reduce the bacterial growth and multiplication with further undesirable changes.

6. REFERENCES

1. Edris, A. A.; Amin-Reham, A.; Naseif- Marionette, Z. and AbdelFatah- Ebtsam, M. (2014): Evaluation of Retailed Salted Fish according to Egyptian Standard. *Benha Veterinary Medical Journal*. 27;168-176.
2. El-Moselhy, K. M. (2000): Accumulation of copper, cadmium and lead in some fish from the Guif of suez. *J Aquat Biol Fish*, 3 (1), 73-83
3. 235–249 Elshafey- Wesam, S.; Ibrahim- Hemmat, M.; Hassan, M.A. and Maarouf, A. A. (2018): Assessment of Histamine and Putrescine Residues in Fish and Shell Fish. *Benha Veterinary Medical Journal* .34(3): 178-187.
4. FDA (2011): Fish and fishery products, hazards and controls guide (3rd Ed.). Pp. 73-90.
5. Feng, C.; Teuber, S. and Gershwin, M.E. (2016): Histamine (Scombroid) Fish Poisoning. *Clinical reviews in allergy & immunology Journal*. 50(1):64-9.
6. Ferran, M. and Yébenes, M. (2006): Flushing associated with scombroid fish poisoning. *Dermatol Online Journal*. 12(6):15.
7. Harmelin, Y.; Hubiche, T.; Pharaon, M. and Del Giudice, P. (2018): [Three cases of scombroid poisoning]. *Annales de Dermatologie et de Venereologie Journal* .145(1):29-32.
8. Hungerford, J.M. (2010): Scombroid poisoning: a review. *Toxicon Journal*. 56(2):231-43.
9. Lee, Y.; Kung, H.; Lin, C.; Hwang, C.; Lin, C. and Tsai, Y. (2012): Histamine production by *Enterobacter aerogenes* in tuna dumpling stuffing at various storage temperatures. *Food Chemistry Journal*. 131: 405-412.
10. Lehane, L. and Olley, J. (2000): Histamine fish poisoning revisited. *International Journal of Food Microbiology*. 58: 1-37.
11. Nader, Y.M.; ElBahy-Engy, F. and Anter-Dalia, M. (2016): Biogenic Amines in Salted and Smoked Herring Fish. *Global Veterinaria Journal*. 16 (6): 525-529.
12. Önal, A. (2006): Current analytical methods for determination of biogenic amines in foods. *Food Chemistry Journal* 103(4): 1475-1786.
13. Patange, S. B.; Mukundan, M. K.; Ashok Kumar, K. 2005. A simple and rapid method for colorimetric determination of histamine in fish flesh. *Food Control Journal* .16, 465–472.
14. Sagratini, G.; Fernández-Frazón, M.; Berardinis, F.; Font, G.; Vittori, S. and Mañes, J. (2012): Determination of eight un derivatized biogenic amines in fish by solid phase extraction and liquid chromatography tandem mass spectrometry. *Food Chemistry Journal*. 132: 537-543.
15. Surya, T.; Sivaraman, B.; Alamelu, V.; A. Priyatharshini, A.; Arisekar, U. and Sundhar, S. (2019): Rapid Methods for Histamine Detection in Fishery Products. *International Journal of Current Microbiology and Applied Sciences*. 8(3): 2035-2046.
16. Tombelli S. and Mascini, M. (1998): Electrochemical biosensors for biogenic amines: a comparison between different approaches. *Analytica Chimica Acta Journal*. 385(3): 277-284.
17. Visciano, P.; Schirona, M.; Tofalo, R. and Suzzi, G. (2012): Biogenic amine in raw and processed seafood. Department of Food science, University of Teramo, Italy. *Frontiers in Microbiology*. 10:33-89.
18. Walaa (2016): Biogenic Amines as quality indices of imported frozen fish M. V. Sc., Thesis, Fac. Vet. Med., Benha University.
19. Zhai, H.; Yang, X.; Li, L.; Xia, G.; Cen, J.; Huang, H. and Hao, S. (2012): Biogenic amines in commercial fish and fish products sold in southern china. *Food Control Journal*. 25: 303–308.