

Evaluation of Viability and Nuclear Status in Vitrified Mature Buffalo Oocytes

¹Mohamed M.M. El-Sokary, ¹Mohamed M.M. Kandiel, ²Karima Gh.M. Mahmoud,
¹Mahmoud E.A. Abouel-Roos and ¹Gamal A.M. Sosa

¹Department of Theriogenology, Faculty of Veterinary Medicine,
Benha University, Kaliobia, Egypt

²Department of Animal Reproduction and A.I.,
National Research Center, Dokki, Giza, Egypt

Abstract: Vitrification is a common method which is successfully used for cryopreservation of gametes and embryos in several animal species, though subsequent progress is still limited, especially in buffalo. The present study aimed to evaluate the effect of vitrification process on nuclear status in matured buffalo oocytes. Morphology, viability, state of nuclear maturation in meiotic stages and maturation rate (matured oocyte/survived oocyte) were assessed in vitrified matured buffalo oocytes. Immature buffalo oocytes were matured *in-vitro* for 22 hrs. and vitrified at room temperature (25°C) in a mixture of ethylene glycol (EG) and dimethyl sulfoxide (DMSO) with 0.25 ml straw. Vitrification solutions (VS) were VS₁ (1.5MEG+ 1.5MDMSO) and VS₂ (3M EG+ 3M DMSO). Cryoprotectants were added in two steps, with the first step concentration half that of the second (and final) step concentration. After warming, normal oocytes were cultured for further 2 hrs., stained with trypan blue for viability evaluation, then fixed and stained with orcein 1% stain for detection of meiotic stages. Oocytes that reached Telophase I (TI) or Metaphase II (MII) stages were considered matured. Results showed that morphologically normal and viable oocytes were significantly decreased in vitrified oocytes than control. The percentage of oocytes reached TI and MII in un-vitrified (control) group was significantly ($P<0.05$) higher than those vitrified after maturation. The maturation rate was higher in control than vitrified oocytes (76.04 ± 2.29 vs. 66.46 ± 2.25). In conclusion, vitrification is a successful method for cryopreservation of matured buffalo oocytes. However, in spite of higher rates of viability and maturation obtained in current study, it is still lower than un-vitrified oocytes.

Key words: Buffalo • Oocytes • Nuclear status • Vitrification

INTRODUCTION

Cryopreservation is a desirable option for generating a readily available source of unfertilized oocytes for both biological and commercial purposes. Vitrification has been attempted by several workers with variable success in mouse [1], human [2], bovine [3], swine [4], equine [5] and buffalo [6] oocytes. Studies have been performed to investigate the best conditions to preserve viability of vitrified oocytes [7]. Oocytes are the only cells which can form a new individual after fertilization. During maturation, oocytes undergo changes in nuclear status involving existence from diplotene stage of the first meiotic prophase, known as germinal vesicle stage and

progression to the metaphase II stage with extrusion of the first polar body [8]. The observation of the oocytes chromosomes at this stage is more reliable mean for defining the *in vitro* maturation progress [9]. Cryoprotectants have been reported not only to induce a disorganization of the spindle microtubules, which in turn results in chromosomal scattering, but also to disrupt the cortical microfilament network. Disruption of the cytoskeleton might be intrinsic to changes in shape and shrinkage that accompany cryopreservation procedures, which in turn might lead to premature release of cortical granules and zona hardening. Also, rewarming may then lead to membrane fusion, cortical granule enzyme release and premature zona hardening [10].

Corresponding Author: Karima Gh. M. Mahmoud, Department of Animal Reproduction and A.I.,
National Research Center, Postal code: 12622, Dokki, Giza, Egypt.
Tel: +2 01001827716; Fax: +2 02 33370931.

The cryopreservation of oocytes with slow cooling or vitrification protocol was reported to have deleterious effect on viability. Oocyte survival rate after cryopreservation is affected by both morphological and biophysical factors [11]. The main biophysical factors causing cellular disruption during cryopreservation are intracellular ice formation [12] and osmotic injury [13]. These factors could be easily minimized by adequate cryopreservation techniques using proper cryoprotectant and its concentration [14]. So the current study was designed to evaluate the effect of vitrification process on the morphology, survivability and nuclear status in vitrified matured buffalo oocytes.

MATERIALS AND METHODS

Chemicals: Chemicals for *in vitro* maturation including fetal calf serum and tissue culture medium (TCM 199) were obtained from Gibco BRL (Grand Island, New York, USA). Cysteamine, ethylene glycol (EG) and dimethyl sulfoxide (DMSO) were obtained from Sigma Chemicals Company.

Oocyte Recovery and Selection: Buffalo (*Bubalus bubalis*) ovaries, collected from local abattoir (El-Moneeb, Giza, Egypt) within 2 hrs. after slaughtering and transported to the laboratory in physiological saline (0.9% NaCl) containing antibiotic (100 µg/ml streptomycin sulfate and 100 IU/ml penicillin) maintained at 30°C. Ovaries were washed three times in phosphate buffered saline (PBS). Oocytes were aspirated from 2 to 5 mm follicles with a 20-gauge needle attached to a 5-ml syringe containing PBS with 3% bovine serum albumin (fraction V) and antibiotics (100 µg/ml streptomycin sulfate and 100 IU/ml penicillin). Oocytes were screened using stereomicroscope and those with intact layers of cumulus cells and homogenous cytoplasm were selected for *in-vitro* maturation [15].

***In vitro* Maturation (IVM) of Oocytes:** Oocyte maturation was carried out as described by Mahmoud [16]. Briefly, the recovered oocytes were cultured in groups of 10 or 20 in 100 µl droplets of maturation medium (TCM-199) supplemented with 10 % fetal calf serum, 50µM cysteamine and 50 µg/ml gentamycin sulfate. Droplets were covered with mineral oil and pre-incubated for a minimum of 2 hrs. in a humidified 5% CO₂ atmosphere at 38.5°C. The oocytes were placed into the droplets and incubated for 22 hrs. in a humidified 5% CO₂ atmosphere at 38.5°C.

Evaluation of Nuclear Maturation by Fluorescence Microscope: After 22 h IVM, COCs were subjected to removal of the surrounding cumulus cells by gentle pipetting with a fine capillary tube. Oocytes were stained with probidium iodide (100 µg/ml) and examined under a fluorescence microscope for nuclear status. Oocytes that reached TelophaseI (TI) or MetaphaseII (MII) stages were considered matured.

Vitrification and Warming: Oocytes were exposed to two-step addition of cryoprotectants [17]. Briefly, oocytes were exposed to VS₁ (1.5 M EG + 1.5 M DMSO) for 45 sec. and VS₂ (3 M EG + 3 M DMSO) for 25 sec. The holding media was TCM 199 containing 2.5 mM HEPES + 20% fetal calf serum. Oocytes (n=5-10) were immediately loaded in 0.25 ml straws in the middle column of VS₂ and separated by air bubbles, sealed with straw plugs, pre-cooled by exposure to liquid nitrogen (LN₂) vapor for at least 60sec and dipped vertically in LN₂ there after for two months. Straws transferred rapidly in less than 5 sec.; to prevent zona fracture [18]; to a water bath at 35-37°C for 20 sec. for thawing and the expelled oocytes were equilibrated for 5 min. in 0.5 M galactose solution [19] in TCM-199 for one-step dilution to remove the cryoprotectants. Oocytes were washed 4-5 times in fresh washing medium and evaluated 24hrs.

Evaluation of Oocytes Morphology after Vitrification-Thawing: Oocyte normality was directly evaluated by an inverted microscope as it was described formerly [20]. Normal oocytes were verified with their spherical and symmetrical shape and absence of lysis signs, membrane damage, swelling, vacuolization, degeneration or leakage of the cellular contents. Abnormal oocytes were characterized by a ruptured zona pellucida or vitelline membrane or having fragmented cytoplasm with signs of degeneration. The normal oocytes were cultured for further 2 hrs. to complete the maturation.

Assessment of Vitrified-Thawed Oocytes Viability by Trypan Blue: Oocyte viability in control (24h-IVM, un-vitrified) and vitrified (2 h post-thawing) was assessed by staining with 0.4% (w/v) trypan blue and examination under an inverted phase-contrast microscope. Exclusion of dye by COCs is an indicative of viability. Viable oocytes with viable or dead cumulus were considered viable while dead oocytes with viable or dead cumulus were considered dead.

Evaluation of Nuclear Maturation by Orcein Stain:

At the end of the culture period, chromosome slides were prepared according to the procedure described by Tarkowski [21]. Briefly, cumulus cells were removed mechanically by gentle pipetting, oocyte was transferred to 1% hypotonic sodium citrate solution for 10 min. and placed on a microscope slide with a minimal amount of hypotonic solution. Three drops of methanol/glacial acetic acid (3:1) fixative were dropped onto the oocytes before staining with 1% orcein stain. The state of nuclear maturation was determined as described earlier by Mahmoud [22]. Oocytes that reached TI or MII stages were considered matured. TI was defined as completion of separation of two chromosome sets and MII was haploid set of chromosomes after emission of first polar body.

Statistical Analysis: Data were presented as mean \pm S.E., tabulated and analyzed for significance by Student *t*-testing SPSS (ver.14) statistical software. Differences were considered to be significant at $P < 0.05$.

RESULTS

Oocytes with complete or partial cumulus cells were selected, matured *in vitro* for 22hrs and their nuclear status was determined by probidium iodide as shown in Fig. 1. Morphological evaluation of *in-vitro* matured vitrified buffalo oocytes directly after thawing and furtherculture for two hours are shown in Table 1 and Fig. 2. The percentage of normal oocytes significantly ($P < 0.05$) decreased in vitrified oocytes than control. Also, the survivability of oocytes was significantly ($P < 0.001$) decreased with vitrification.

The stages of nuclear maturation in control and vitrified oocytes as determined by orcein staining and microscopic examination are shown in Table 2. Cytogenetic analysis of 209 control and 167 vitrified oocytes showed a wide range of meiotic configurations: Metaphase-I, Anaphase-I, Telophase-I and Metaphase-II (Fig. 3). The percentage of oocytes reached TI and MII in control oocytes was significantly ($P < 0.05$) higher than those vitrified after maturation. The mean maturation rate in control oocytes (76.04 ± 2.29) was higher than oocytes exposed to vitrification (66.46 ± 2.25).

Table 1: Morphological evaluation and survivability directly after thawing and two hours culture of vitrified mature buffalo oocytes

Oocytes culture	No. oocytes examined	Morphologically. Normal	Survivability 2hrs. post-thawing
		No %	No %
Vitrified	255	226 (89.33 \pm 1.8)	170 (75.29 \pm 0.8)
Control (un-vitrified)	245	242 (98.60 \pm 0.7)*	214 (88.50 \pm 0.4)**

(* $P < 0.05$ -** $P < 0.001$)

Table 2: Effect of vitrification on nuclear status of *in-vitro* matured buffalo oocytes

Oocyte type	n	Stage of nuclear maturation						Maturation rate (TI+MII) No. (mean \pm S.E.)
		Germinal vesicle No. (%)	GVBD No. (%)	Metaphase I (MI) No. (%)	Anaphase I (AI) No. (%)	Metaphase II (MII) No. (%)	Telophase I (TI) No. (%)	
Control	209	3 (15.6)	5 (2.7)	5 (2.7)	6 (2.9)	123 (58.4)	37 (17.4)	160(76.04 \pm 2.29)*
Vitrified	167	0 (13.1)	5 (10.1)	6 (3.2)	9 (5.7)	82 (48.2)	35 (20.6)	117 (66.46 \pm 2.25)

n: number of oocytes, GVBD: Germinal Vesicle Break Down. Maturation rate (matured oocyte (TI+MII)/survived oocyte) with superscript (*) indicated a significant difference between verified and control oocytes at $P < 0.05$.




Fig. 1: Buffalo oocytes at Telophase I (A) and Metaphase II (B) stages stained with probidium iodide and visualized under fluorescence microscope ($\times 400$).




Fig. 2: Morphological evaluation of *in-vitro* matured vitrified buffalo oocytes stained by trypan blue. (A) viable oocyte and viable cumulus cells, (B) viable oocyte and non-viable cumulus cells, (C) non-viable oocyte and viable cumulus cells and (D) non-viable oocyte and non-viable cumulus cells.




Fig. 3: Buffalo oocytes at germinal vesicle break down with a condensation of chromatin (A), Metaphase I stage (B), anaphase I stage with complete homologous segregation of chromosomes (C), Telophase I stage with two groups of equally spread homologous chromosomes (D), Metaphase II. Note the normal haploid number and the first polar body chromosomes have undergone degeneration (E). Metaphase II stage (F).

DISCUSSION

Oocytes meiosis is very sensitive to endogenous and exogenous factors that could result in oocytes with chromosomal abnormalities. They might be affected by temperature reduction, cryoprotectant concentration and *in vitro* culture [23]. Cytogenetic analysis in the present study showed a wide range of meiotic configurations in Metaphase I, Anaphase I, Telophase I and Metaphase II. During maturation, oocytes undergo changes in nuclear status that involve exit from diplotene stage of the first meiotic prophase, known as germinal vesicle stage and progression to the metaphase II stage with extrusion of the first polar body [8]. The observation of the oocytes chromosomes at this stage is more reliable mean for defining the *in vitro* maturation progress [9].

The survival rate of vitrified-warmed oocytes observed in the present study was low as compared with control. This in accordance with data reported previously in buffalo oocytes [17, 24-26]. In fact, all oocytes and embryos suffer considerable morphological and functional damage during cryopreservation, but the extent of the injury as well as differences in post-thaw survival and developmental rates may be highly variable depending on the species, developmental stage and origin. Oocytes are particularly difficult to be cryopreserved successfully and thus resulted in low rates of blastocyst production after thawing, fertilization and culture [28]. This problem may be due to the large size of oocytes which consequently have a low surface to volume ratio, making it more difficult for water and cryoprotectants to move across the cell plasma membranes.

In the current work, maturation rate; represented by the percentage of oocytes reached TI and MII; decreased in vitrified than control group. It is well known that vitrification causes several ultra-structural [28] and structural alterations, including damages of the meiotic spindle apparatus in the oocytes [29]. Disorganization of meiotic spindles might lead to chromosomal dispersion, aneuploidy or polyploidy of *in-vitro* matured cryopreserved oocytes, as well as failure of normal fertilization and development [30]. The evidence of these structural alterations provides the basis for developing new strategies to improve the efficiency of cryopreservation in cattle and buffalo species [31].

In conclusion, vitrification is a successful method for cryopreservation of matured buffalo oocytes but still has adverse effects on the viability and percentage of oocytes reached Metaphase II.

REFERENCES

1. Edashige, K., Y. Yamaji, F.W. Kleinhans and M. Kasai, 2003. Artificial expression of Aquaporin-3 improves the survival of mouse oocytes after cryopreservation. *Biol. Reprod.*, 68: 87-94.
2. Katayama, K.P., J. Stehlik, M. Kuwayana, O. Kato and E. Stehlik, 2003. High survival rate of vitrified human oocytes results in clinical pregnancy. *Fert. Steril.*, 80: 223-224.
3. Albarracin, J.L., R. Morato, D. Izquierdo and T. Mogas, 2005. Vitrification of calf oocytes: Effects of maturation stage and prematuration treatment on the nuclear and cytoskeletal components of oocytes and their subsequent development. *Mol. Reprod. Dev.*, 72: 239-249.
4. Huang, W.T. and W. Holtz, 2002. Effects of meiotic stages, cryoprotectants, cooling and vitrification on cryopreservation of porcine oocytes. *Asian-Aust. J. Anim. Sci.*, 15: 485-493.
5. Hurtt, A.E., F. Landim-Alvarenga, G.E. Seidel and E.L. Squires, 2000. Vitrification of immature and mature equine and bovine oocytes in ethylene glycol, Ficoll and sucrose solution using open-pulled straws. *Theriogenology*, 54: 119-128.
6. Hufana-Duran, D., P.B. Pedro, H.V. Venturina, R.D. Hufana, A.L. Salazar, P.G. Duran and L.C. Cruz, 2004. Post-warming hatching and birth of live calves following transfer of *in vitro*-derived vitrified water buffalo. *Theriogenology*, 61: 1429-1439.
7. Vajta, G. and M. Kuwayama, 2006. Improving cryopreservation systems. *Theriogenology*, 65: 236-244.
8. Edwards, R.G., 1962. Meiosis in ovarian oocytes of adult mammals. *J. Nature*, 196: 446-450.
9. Mahmoud, K. Gh. M. and N.T. Eashra, 2004. Meiotic chromosomes of cattle oocytes in relation to seasonal variation. *Egypt. J. Vet. Sci.*, 38: 1-9.
10. Vincent, C. and M.H. Johnson, 1992. Cooling, cryoprotectants and the cytoskeleton of the mammalian oocyte. *Oxf. Rev. Reprod. Biol.*, 14: 73-100.
11. Ledda, S., L. Bogliolo, S. Succu, F. Ariu, D. Bebbere, G.G. Leoni and S. Naitana, 2007. Oocyte cryopreservation: oocyte assessment and strategies for improving survival. *Reprod. Fertil. Dev.*, 19: 13-23.
12. Mazur, P., S. Seki, I.L. Pinn, F.W. Kleinhans and K. Edashige, 2005. Extra- and intracellular ice formation in mouse oocytes. *Cryobiology*, 51: 29-53.
13. Pedro, P.B., S.E. Zhu, N. Makino, T. Sakurai, K. Edashige and M. Kasai, 1997. Effects of hypotonic stress on the survival of mouse oocytes and embryos at various stages. *Cryobiology*, 35: 150-158.
14. Wani, N.A., S.N. Maurya, A.K. Misra, V.B. Saxena and B.D. Lakhchaura, XXXX. Effect of cryoprotectants and their concentration on *in vitro* development of vitrified warmed immature oocytes in buffalo (*Bubalus bubalis*). *Theriogenology*, 61: 831-842.
15. Warriach, H.M. and K.R. Chohan, 2004. Thickness of cumulus cell layer is a significant factor in meiotic competence of buffalo oocytes. *J. Vet. Sci.*, 5: 247-251.
16. Mahmoud, K. Gh. M., 2001. Cytogenetic studies on *in-vitro* fertilization in buffaloes. Ph.D. Thesis, Fac. Vet. Med., Cairo University, Egypt.
17. Mahmoud, K. Gh. M., T.H. Scholkamy, Y.F. Ahmed, Jr. G.E. Seidel and M.F. Nawito, 2010a. Effect of different combinations of cryoprotectants on *in vitro* maturation of immature buffalo (*Bubalus bubalis*) oocytes vitrified by straw and open pulled straw methods. *Reprod. Dom. Anim.*, 45: 565-571.
18. Subramaniam, A., K.P. Devarajan and M. Mohanan, 1990. Incidence and avoidance of zona fracture in cryopreserved bovine ova and embryos. *N.Z. Vet. J.*, 38: 156-157.
19. Horvath, G. and G. E.Jr. Seidel, 2006. Vitrification of bovine oocytes after treatment with cholesterol-loaded methyl- β -cyclodextrin. *Theriogenology*, 66: 1026-1033.
20. Dhali, A., R.S. Manik, S.K. Das, S.K. Singla and P. Palta, 2000. Effect of ethylene glycol concentration and exposure time on post-vitrification survival and *in vitro* maturation rate of buffalo oocytes. *Theriogenology*, 50: 521-530.

21. Tarkowski, A.K., 1966. An air-drying method for chromosome preparations from mouse eggs. *Cytogenetics*, 5: 394-400.
22. Mahmoud, K.Gh.M., 2004. Meiotic stages and incidence of diploid oocytes in Egyptian cattle and buffaloes. *Assiut Vet. Med. J.*, 50: 336-347.
23. Van der Elst, J., E. Van den Abbeel, S. Nerinckx and A. Van Steirteghem, 1992. Parthenogenetic activation pattern and microtubular organization of the mouse oocyte after exposure to 1, 2 propandiol. *Cryobiology*, 29: 549-562.
24. Mahmoud, K.Gh.M., T.H. Scholkamy, Jr. G.E. Seidel and G.A. Elsisy, 2010b. Influence of Meiotic stages on the survival and developmental ability of vitrified-warmed buffalo oocytes. *GJBRR*, 5: 220-225.
25. Liang, Y.Y., K. Srirattana, T. Phermthai, T. Somfai, T. Nagai and R. Parnpai, 2012. Effects of vitrification cryoprotectant treatment and cooling method on the viability and development of buffalo oocytes after intracytoplasmic sperm injection. *Cryobiology*, 65: 151-156.
26. Attanasio, L., L. Boccia, G. Vajta, M. Kuwayama, G. Campanile, L. Zicarelli, G. Neglia and B. Gasparrini, 2010. Cryotop vitrification of buffalo (*Bubalus bubalis*) *in vitro* matured oocytes: effects of cryoprotectant concentrations and warming procedures. *Reprod. Domest. Anim.*, 45: 997-1002.
27. Vajta, G., P. Holm, M. Kuwayama, P.J. Booth, H. Jacobsen, T. and Greve and H. Callesen, 1998. Open pulled straw (OPS) vitrification: a new way to reduce cryoinjuries of bovine ova and embryos. *Mol.Reprod. Dev.*, 51: 53-58.
28. Hyttel, P., G. Vajta and H. Callesen, 2000. Vitrification of bovine oocytes with the open pulled straw method: ultrastructural consequences. *Mol. Reprod. Dev.*, 56: 80-88.
29. Succu, S., D. Bebbere, L. Bogliolo, F. Ariu, S. Fois, G.G. Leoni, F. Berlinguer, S. Naitana and S. Ledda, 2008. Vitrification of *in vitro* matured ovine oocytes affects *in vitro* pre-implantation development and mRNA abundance. *Mol. Reprod. Dev.*, 75: 538-546.
30. Chen, S.U., Y.R. Lien, K.H. Chao, H.N. Ho, Y.S. Yang and T.Y. Lee, 2003. Effects of cryopreservation on meiotic spindles of oocytes and its dynamics after thawing: clinical implications in oocyte freezing. A review. *Mol. Cell. Endocrinol.*, 202: 101-107.
31. Gasparrini, B., S. Francesco and M. Rubessa, 2009. Structural changes of *in vitro* matured buffalo and bovine oocytes following cryopreservation. *Ital. J. Anim. Sci.*, 8: 90-92.